

ON THE MINIMA AND CONVEXITY OF EPSTEIN ZETA FUNCTION

S.C. LIM¹ AND L.P. TEO²

¹Faculty of Engineering, Multimedia University, Jalan Multimedia,
Cyberjaya, 63100, Selangor Darul Ehsan, Malaysia.

²Faculty of Information Technology, Multimedia University, Jalan Multimedia,
Cyberjaya, 63100, Selangor Darul Ehsan, Malaysia.

ABSTRACT. Let $Z_n(s; a_1, \dots, a_n)$ be the Epstein zeta function defined as the meromorphic continuation of the function

$$\sum_{k \in \mathbb{Z}^n \setminus \{0\}} \left(\sum_{i=1}^n [a_i k_i]^2 \right)^{-s}, \quad \text{Re } s > \frac{n}{2}$$

to the complex plane. We show that for fixed $s \neq n/2$, the function $Z_n(s; a_1, \dots, a_n)$, as a function of $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$ with fixed $\prod_{i=1}^n a_i$, has a unique minimum at the point $a_1 = \dots = a_n$. When $\sum_{i=1}^n c_i$ is fixed, the function

$$(c_1, \dots, c_n) \mapsto Z_n(s; e^{c_1}, \dots, e^{c_n})$$

can be shown to be a convex function of any $(n-1)$ of the variables $\{c_1, \dots, c_n\}$. These results are then applied to the study of the sign of $Z_n(s; a_1, \dots, a_n)$ when s is in the critical range $(0, n/2)$. It is shown that when $1 \leq n \leq 9$, $Z_n(s; a_1, \dots, a_n)$ as a function of $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$, can be both positive and negative for every $s \in (0, n/2)$. When $n \geq 10$, there are some open subsets $I_{n,+}$ of $s \in (0, n/2)$, where $Z_n(s; a_1, \dots, a_n)$ is positive for all $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$. By regarding $Z_n(s; a_1, \dots, a_n)$ as a function of s , we find that when $n \geq 10$, the generalized Riemann hypothesis is false for all $(a_1, \dots, a_n)$.

1. INTRODUCTION

In [1, 2], Epstein introduced the following two-dimensional zeta function

$$Z_2(A; s) = \sum_{(j,k) \in \mathbb{Z}^2 \setminus \{(0,0)\}} [A(j,k)]^{-s}, \quad \text{Re } s > 1,$$

where

$$A(j,k) = aj^2 + 2bjk + ck^2 = \begin{pmatrix} j & k \end{pmatrix} \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} j \\ k \end{pmatrix}$$

is a positive definite real quadratic form associated to the positive definite symmetric matrix

$$A = \begin{pmatrix} a & b \\ b & c \end{pmatrix}.$$

2000 *Mathematics Subject Classification.* Primary 11E45, 26B15 .

Key words and phrases. Epstein Zeta function, convexity, minima.

Later on, some generalizations of this zeta function to higher dimension were considered [3, 4, 5, 6, 7]. One of the generalizations is given by

$$(1.1) \quad Z_n(A; s) = \sum_{k \in \mathbb{Z}^n \setminus \{0\}} [k^t A k]^{-s}, \quad \operatorname{Re} s > \frac{n}{2},$$

where now A is an $n \times n$ positive definite symmetric matrix and

$$k^t A k = \sum_{i=1}^n \sum_{j=1}^n A_{ij} k_i k_j$$

is the associated quadratic form. It was proved that the Epstein zeta function $Z_n(A, s)$ has a meromorphic continuation to the whole complex plane with a single pole at $s = n/2$, and it satisfies a functional equation. When $n = 1$ and $A = (1)$, $Z_1(A; s)$ is nothing but equal to $2\zeta_R(2s)$, where $\zeta_R(s)$ is the Riemann zeta function. The generalized Riemann hypothesis associated to the Epstein zeta function says that aside from the trivial zeros at $s = -j, j \in \mathbb{N}$, all the other zeros of $Z_n(A; s)$ are located at the critical line $\operatorname{Re} s = \frac{n}{4}$. As far back as 1947, it has been known [8] that the Riemann hypothesis is in general not true for the Epstein zeta function, even for $n = 2$.

There exists another interpretation of Epstein zeta function with more geometric flavour. Let $v_1, \dots, v_n$ be n linearly independent vectors in $\mathbb{R}^n$ and let L be the non-degenerate lattice generated by $\{v_1, \dots, v_n\}$, i.e.

$$L = \left\{ \sum_{i=1}^n k_i v_i : (k_1, \dots, k_n) \in \mathbb{Z}^n \right\}.$$

The quotient of $\mathbb{R}^n$ by L , i.e. $\mathbb{R}^n/L$, is an n -dimensional torus. The function

$$\zeta_n(L; s) = \sum_{v \in L \setminus \{0\}} \langle v, v \rangle^{-s}, \quad \text{where } \langle v, w \rangle := \sum_{i=1}^n v_i w_i,$$

is called the zeta function associated to the lattice L . It is actually the same as the Epstein zeta function $Z_n(A; s)$ if $A = T_L^t T_L$, and T_L being the $n \times n$ matrix whose columns are the generators $v_1, \dots, v_n$ of the lattice L , i.e.,

$$T_L = (v_1 \quad \dots \quad v_n).$$

Given a lattice L , its dual lattice L^* is defined as the set

$$L^* = \left\{ w \in \mathbb{R}^n : \langle w, v \rangle = \sum_{i=1}^n w_i v_i \in \mathbb{Z} \text{ for all } v \in L \right\}.$$

L^* is a lattice generated by a dual set of vectors $\{w_1, \dots, w_n\}$ with $\langle w_i, v_j \rangle = \delta_{ij}$. The corresponding matrix T_{L^*} is related to T_L by $T_{L^*} = (T_L^{-1})^t$. The zeta function $\zeta_n(L^*; s)$ of the dual lattice L^* is related to a spectral zeta function associated to the compact torus $\mathbb{R}^n/L$. Given a compact manifold M with Laplace operator Δ_M , the zeta function of Δ_M is defined as

$$(1.2) \quad \zeta_{\Delta_M}(s) = \sum_{n=1}^{\infty} \lambda_n^{-s},$$

where $0 < \lambda_1 \leq \lambda_2 \leq \dots$ are the nonzero eigenvalues of Δ_M . For the compact torus $\mathbb{R}^n/L$, the Laplace operator $\Delta = \frac{\partial^2}{\partial x_1^2} + \dots + \frac{\partial^2}{\partial x_n^2}$ on $\mathbb{R}^n$ descends to the Laplace

operator on $\mathbb{R}^n/L$. One can easily check that the associated zeta function $\zeta_{\Delta_{\mathbb{R}^n/L}}$ is up to a constant, the zeta function of the dual lattice L^* , i.e.

$$\zeta_{\Delta_{\mathbb{R}^n/L}} = c\zeta_n(L^*; s) = cZ_n(T_L^{-1}(T_L^{-1})^t; s).$$

Under this perspective, the Epstein zeta function enters the realm of theoretical physics. In quantum theory, very often one needs to compute the functional determinant of a positive definite (pseudo)-differential operator P (e.g., the Laplace operator), which is defined as

$$(1.3) \quad \mathfrak{D}_P = \prod_{i=1}^{\infty} \lambda_i,$$

where $\lambda_i, 1 \leq i < \infty$ are the nonzero eigenvalues of P . This functional determinant usually appears as one-loop partition function of a field theory, and is closely related to Casimir effect as well as the one-loop effective potential of the theory. In general, the infinite product in (1.3) is divergent and regularization is required to obtain a finite quantity. One of the regularization techniques, known as zeta regularization method (see e.g. [9, 10, 11]), uses the zeta function associated to P , $\zeta_P(s)$, which is defined as in (1.2). By proving that $\zeta_P(s)$ has an analytic continuation to a neighborhood of $s = 0$, the regularized functional determinant $\mathfrak{D}_P$ is then defined as

$$\mathfrak{D}_P = \exp(-\zeta'_P(0)).$$

When the underlying spacetime of the field theory is a toroidal manifold $T^n \times \mathbb{R}^N$, where T^n is an n -dimensional torus with compactification lengths $L_1, \dots, L_n$ and P is the Laplace operator, the associated zeta function $\zeta_P(s)$ is, up to a constant, equal to

$$\sum_{k \in \mathbb{Z}^n \setminus \{0\}} \left(\sum_{i=1}^n \left[\frac{k_i}{L_i} \right]^2 \right)^{-s + \frac{N}{2}},$$

which is the Epstein zeta function $Z_n(A; s - \frac{N}{2})$, with $A = \text{diag}\{1/L_1^2, \dots, 1/L_n^2\}$ a diagonal matrix. We use the notation

$$(1.4) \quad Z_n(s; a_1, \dots, a_n) := \sum_{k \in \mathbb{Z}^n \setminus \{0\}} \left(\sum_{i=1}^n [a_i k_i]^2 \right)^{-s}$$

to denote this special subclass of the Epstein zeta function corresponding to

$$A = \begin{pmatrix} a_1^2 & 0 & \dots & 0 \\ 0 & a_2^2 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & a_n^2 \end{pmatrix}$$

in (1.1). Here it is assumed that $a_1, \dots, a_n$ are positive real numbers. Another situation where $Z_n(s; a_1, \dots, a_n)$ appears is when the underlying spacetime is a rectangular cavity $\Omega := [0, L_1] \times \dots \times [0, L_n] \times \mathbb{R}^N$. The function $Z_n(s; a_1, \dots, a_n)$ satisfies a functional equation (or called reflection formula), which can be written

in the following symmetric form

$$(1.5) \quad \sqrt{\prod_{i=1}^n a_i} \pi^{-s} \Gamma(s) Z_n(s; a_1, \dots, a_n) = \frac{\pi^{s-\frac{n}{2}}}{\sqrt{\prod_{i=1}^n a_i}} \Gamma\left(\frac{n}{2} - s\right) Z_n\left(\frac{n}{2} - s; \frac{1}{a_1}, \dots, \frac{1}{a_n}\right).$$

In physics literature, the Epstein zeta function $Z_n(s; A)$ usually appears together with gamma function in the combination

$$(1.6) \quad \Gamma(s) Z_n(s; A).$$

For example, in [12], it is found that the Casimir energy of a massless scalar field in the rectangular cavity $\Omega = [0, L_1] \times \dots \times [0, L_n] \times \mathbb{R}^N$ under periodic boundary conditions is, up to a positive constant, equal to

$$(1.7) \quad -\Gamma\left(-\frac{N}{2}\right) Z_n\left(-\frac{N}{2}; \frac{1}{L_1}, \dots, \frac{1}{L_n}\right).$$

For massless scalar fields under Dirichlet or Neumann boundary conditions, or electromagnetic fields in cavities with perfectly conducting walls or walls with infinite permeability, the corresponding Casimir energy is expressible as linear combinations of expressions of the form (1.7). Another example is the massless scalar field theory with $\lambda\varphi^4$ interaction on the toroidal manifold $T^n \times \mathbb{R}^{4-n}$, where $n = 1, 3, 4$. It was found that [13, 14, 15, 16, 17] the topologically generated mass of this theory, to the one-loop order, is, up to a positive constant, given by :

$$\Gamma\left(\frac{n}{2} - 1\right) Z_n\left(\frac{n}{2} - 1; \frac{1}{L_1}, \dots, \frac{1}{L_n}\right).$$

In view of the functional equation (1.5) and the importance of the combination (1.6), we define the Xi-function as

$$(1.8) \quad \Xi_n(s; a_1, \dots, a_n) = \sqrt{\prod_{i=1}^n a_i} \pi^{-s} \Gamma(s) Z_n(s; a_1, \dots, a_n).$$

In the study of Casimir effect, the attractive or repulsive nature of the Casimir force, and the geometric configuration of the spacetime that will minimize or maximize the Casimir effect are important issues. In the study of interacting scalar field theory, the sign of the topologically generated mass determines whether symmetry breaking mechanism occurs. Therefore it is important to study the sign of the Xi-function (1.8) and determine its extrema as a function of $(a_1, \dots, a_n)$. This paper is devoted to the study of these issues. In fact, the main motivation comes from our recent study of $\lambda\varphi^4$ interacting fractional Klein–Gordon field theory on a toroidal manifold $T^n \times \mathbb{R}^N$ [18]. In that study, we find that to the one-loop order, the topological generated mass of a α -fractional massless Klein–Gordon field is given by

$$\begin{aligned} m_T^2 &= \frac{\lambda}{\Gamma(\alpha)} \frac{1}{2^{2\alpha+1} \pi^{2\alpha-\frac{N}{2}} [\prod_{i=1}^n L_i]} \Gamma\left(\alpha - \frac{N}{2}\right) Z_n\left(\alpha - \frac{N}{2}; \frac{1}{L_1}, \dots, \frac{1}{L_n}\right) \\ &= \frac{\lambda}{\Gamma(\alpha)} \frac{1}{2^{2\alpha+1} \pi^{\frac{n+N}{2}}} \Gamma\left(\frac{n+N}{2} - \alpha\right) Z_n\left(\frac{n+N}{2} - \alpha; L_1, \dots, L_n\right) \end{aligned}$$

when $\alpha \neq N/2$ or $(n+N)/2$. We need to study the sign of m_T^2 as a function of α, n, N and $L_1, \dots, L_n$ to determine whether symmetry breaking appears. To solve

this problem, it is necessary to determine the minimum of $\Xi_n(s; L_1, \dots, L_n)$ for fixed $\prod_{i=1}^n L_i$ and s .

The study of the (local) minima of the general Epstein zeta function $Z_n(A; s)$ (1.1) as a function of $A \in \{\text{symmetric positive definite } n \times n \text{ real matrices}\} = \mathcal{P}_n$ with fixed determinant has a long history [15–38]. Such investigation is important from a number of points of view. One can read the introduction of the recent paper [42] for an overview of this problem. The local minima of $Z_n(A; s)$, $A \in \mathcal{P}_n$, $\det A = 1$, for s in some domain of $\mathbb{R}$ has only been determined for $n = 2, 3, 4, 5, 6, 7, 8, 24$; and the problem is far from being solved. In this paper, we restrict ourselves to a milder problem of determining the minimum of $Z_n(A; s)$ in the subspace of positive definite symmetric forms consists of diagonal forms. We give an elementary proof to show that for any n and s , if $\prod_{i=1}^n a_i$ is fixed, the minimum of $\Xi_n(s; a_1, \dots, a_n)$ (and hence of $Z_n(s; a_1, \dots, a_n)$) appears at the point $a_1 = \dots = a_n$. This result implies that if $\Xi_n(s) = \Xi_n(s; 1, \dots, 1) \geq 0$, then $\Xi_n(s; a_1, \dots, a_n) \geq 0$ for all $(a_1, \dots, a_n)$. Therefore, the problem whether $\Xi_n(s; a_1, \dots, a_n)$ will be negative for some $(a_1, \dots, a_n)$ is reduced to the problem of whether $\Xi_n(s)$ is negative. In section 3, we give a detail study of the sign of the function $\Xi_n(s)$. We show that for $1 \leq n \leq 9$, $\Xi_n(s)$ is negative for all $s \in (0, n/2)$; whereas for $n \geq 10$, there is a nonempty subset of $(0, n/2)$ where $\Xi_n(s)$ is positive. In section 4, we prove that for fixed s , when $\sum_{i=1}^n c_i$ is fixed, the function

$$(1.9) \quad (c_1, \dots, c_n) \mapsto Z_n(s; e^{c_1}, \dots, e^{c_n})$$

is a convex function of any $(n-1)$ of the variables $\{c_1, \dots, c_n\}$. In fact, we prove a stronger result. We show that for a function of the form

$$\mathcal{F}_n(x_1, \dots, x_n) = \prod_{i=1}^n f(x_i),$$

where $f : \mathbb{R} \rightarrow \mathbb{R}^+$ is a positive twice continuously differentiable function, $\mathcal{F}_n$ is convex if the function $\log f(x)$ is convex. Finally, the result about the convexity of the function (1.9) can be used to obtain conclusion on the connectivity and convexity of some regions of $(c_1, \dots, c_n)$ where $Z_n(s; e^{c_1}, \dots, e^{c_n})$ is negative.

2. THE MINIMUM OF EPSTEIN ZETA FUNCTION

In this section, we show that for $s \notin \{n/2\} \cup \{0, -1, -2, \dots\}$, the function $Z_n(s; a_1, \dots, a_n)$, $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$ has a minimum at $a_1 = \dots = a_n$ when $\prod_{i=1}^n a_i$ is fixed.

There are various ways to obtain the meromorphic continuation of the Epstein zeta function $Z_n(s; a_1, \dots, a_n)$ (1.4). Two of them will be useful to us. The first one has the form (see e.g. [42]):

$$(2.1) \quad \begin{aligned} V \pi^{-s} \Gamma(s) Z_n(s; a_1, \dots, a_n) &= \Xi_n(s; a_1, \dots, a_n) \\ &= -\frac{V}{s} - \frac{V^{-1}}{\left(\frac{n}{2} - s\right)} + V \int_1^\infty t^{s-1} \sum_{k \in \mathbb{Z}^n \setminus \{0\}} \exp\left(-\pi t \sum_{i=1}^n [a_i k_i]^2\right) dt \\ &\quad + V^{-1} \int_1^\infty t^{\frac{n}{2}-s-1} \sum_{k \in \mathbb{Z}^n \setminus \{0\}} \exp\left(-\pi t \sum_{i=1}^n \left[\frac{k_i}{a_i}\right]^2\right) dt, \end{aligned}$$

where

$$V = \sqrt{\prod_{i=1}^n a_i}.$$

This formula shows clearly the Xi-function $\Xi_n(s; a_1, \dots, a_n)$ only has simple poles at $s = 0$ and $s = \frac{n}{2}$. Moreover, since the gamma function $\Gamma(s)$ has simple poles at $s = 0, -1, -2, -3, \dots$, the Epstein zeta function $Z_n(s; a_1, \dots, a_n)$ is identically zero at the negative integer points $-1, -2, -3, \dots$. The second formula is one form of the Chowla-Selberg formula [43, 44]:

$$(2.2) \quad \pi^{-s} \Gamma(s) Z_n(s; a_1, \dots, a_n) = 2a_1^{-2s} \pi^{-s} \Gamma(s) \zeta_R(2s) + 2 \sum_{j=1}^{n-1} \frac{\pi^{-s+\frac{j}{2}} \Gamma(s - \frac{j}{2})}{a_{j+1}^{2s-j} \prod_{l=1}^j a_l} \zeta_R(2s-j) + 4 \sum_{j=1}^{n-1} \frac{1}{\prod_{l=1}^j a_l} \times \\ \sum_{\mathbf{k} \in \mathbb{Z}^j \setminus \{0\}} \sum_{p=1}^{\infty} \frac{1}{(pa_{j+1})^{s-\frac{j}{2}}} \left(\sum_{l=1}^j \left[\frac{k_l}{a_l} \right]^2 \right)^{\frac{s-j}{2}-\frac{j}{4}} K_{s-\frac{j}{2}} \left(2\pi p a_{j+1} \sqrt{\sum_{l=1}^j \left[\frac{k_l}{a_l} \right]^2} \right),$$

where $\zeta_R(s) = \sum_{k=1}^{\infty} k^{-s}$ is the Riemann zeta function. From (2.2), it is easy to verify that for any $\lambda \in \mathbb{R}^+$,

$$Z_n(s; \lambda a_1, \dots, \lambda a_n) = \lambda^{-2s} Z_n(s; a_1, \dots, a_n).$$

Therefore, it only makes sense to look for the minimum of $Z_n(s; a_1, \dots, a_n)$ as a function of $(a_1, \dots, a_n)$ when V is fixed. Without loss of generality, it suffices to consider $V = 1$. We are going to make use of (2.1) to find the minimum of the Xi-function $\Xi_n(s; a_1, \dots, a_n)$ as a function of $(a_1, \dots, a_n)$ for fixed $s \neq 0, n/2$ and when $V = 1$. Since multiplying a constant does not affect the minimum of a function, this will give the minimum of the Epstein zeta function $Z_n(s; a_1, \dots, a_n)$ as a function of $(a_1, \dots, a_n)$ for $s \notin \{n/2\} \cup \{0, -1, -2, \dots\}$ and when $V = 1$. By (2.1), for fixed s and $V = 1$, the minimum of $\Xi_n(s; a_1, \dots, a_n)$ is the same as the minimum of

$$(2.3) \quad \Lambda_n(s; a_1, \dots, a_n) = \int_1^{\infty} t^{s-1} \sum_{k \in \mathbb{Z}^n \setminus \{0\}} \exp \left(-\pi t \sum_{i=1}^n [a_i k_i]^2 \right) dt \\ + \int_1^{\infty} t^{\frac{n}{2}-s-1} \sum_{k \in \mathbb{Z}^n \setminus \{0\}} \exp \left(-\pi t \sum_{i=1}^n \left[\frac{k_i}{a_i} \right]^2 \right) dt.$$

Define the theta function $\vartheta(t)$ by

$$(2.4) \quad \vartheta(t) = \vartheta_3(0, e^{-\pi t}) = \sum_{k=-\infty}^{\infty} e^{-\pi t k^2}$$

Here $\vartheta_3(z, q)$ is a Jacobi theta function [45]. The theta function $\vartheta(t)$ satisfies a reflection formula:

$$(2.5) \quad \vartheta(t) = \frac{1}{\sqrt{t}} \vartheta \left(\frac{1}{t} \right).$$

The theta function (2.4) can be used to reexpress (2.3) as

$$(2.6) \quad \Lambda_n(s; a_1, \dots, a_n) = \int_1^\infty t^{s-1} \left(\prod_{i=1}^n \vartheta(ta_i^2) - 1 \right) dt \\ + \int_1^\infty t^{\frac{n}{2}-s-1} \left(\prod_{i=1}^n \vartheta\left(\frac{t}{a_i^2}\right) - 1 \right) dt.$$

To determine the minimum of $\Lambda_n(s; a_1, \dots, a_n)$, we need the following:

Proposition 2.1. *The function $\log \vartheta(e^u)$, $u \in \mathbb{R}$ is strictly convex. Namely, for any m distinct points $u_1, \dots, u_m \in \mathbb{R}$ and m constants $\lambda_1, \dots, \lambda_m \in (0, 1)$ such that $\sum_{i=1}^m \lambda_i = 1$, we have*

$$\log \vartheta \left(\exp \left[\sum_{i=1}^m \lambda_i u_i \right] \right) < \sum_{i=1}^m \lambda_i \log \vartheta(e^{u_i}).$$

This proposition can be used to prove the main result of this section.

Theorem 2.2. *For fixed $s \in \mathbb{R} \setminus \{0, n/2\}$, the function $\Xi_n(s; a_1, \dots, a_n)$, where $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$ with $\prod_{i=1}^n a_i = 1$, has a unique minimum at $a_1 = \dots = a_n = 1$. Therefore for fixed $s \in \mathbb{R} \setminus (\{n/2\} \cup \{0, -1, -2, -3, \dots\})$, the same statement holds for the function $Z_n(s; a_1, \dots, a_n)$.*

First we show how we prove this theorem from Proposition 2.1.

Proof. By the strict convexity of $\log \vartheta(e^u)$ asserted in Proposition 2.1, we find that if $(a_1, \dots, a_n) \neq (1, \dots, 1)$, then

$$\log \prod_{i=1}^n \vartheta(a_i^2 t) = \sum_{i=1}^n \log \vartheta(\exp[\log t + \log a_i^2]) \\ > n \log \vartheta \left(\exp \left\{ \frac{1}{n} \sum_{i=1}^n [\log t + \log a_i^2] \right\} \right) = \log \vartheta(t)^n.$$

Therefore,

$$(2.7) \quad \prod_{i=1}^n \vartheta(a_i^2 t) > \vartheta(t)^n.$$

Similarly,

$$(2.8) \quad \prod_{i=1}^n \vartheta\left(\frac{t}{a_i^2}\right) > \vartheta(t)^n.$$

Since $t^{\beta-1} > 0$ for all $t > 0$ and $\beta \in \mathbb{R}$ and

$$\prod_{i=1}^n \vartheta(b_i^2 t) - 1 > 0 \quad \text{for all } (b_1, \dots, b_n) \in (\mathbb{R}^+)^n,$$

we conclude from (2.6), (2.7) and (2.8) that for all $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n \setminus \{(1, \dots, 1)\}$ with $\prod_{i=1}^n a_i = 1$, the inequality

$$\Lambda_n(s; a_1, \dots, a_n) > \Lambda_n(s; 1, \dots, 1)$$

holds. The assertion of the theorem follows. $\square$

Now we return to the proof of Proposition 2.1.

Proof. Let

$$(2.9) \quad \begin{aligned} g(u) &= \log \vartheta(u), & u \in \mathbb{R}^+, \\ f(u) &= g(e^u) = \log \vartheta(e^u), & u \in \mathbb{R}. \end{aligned}$$

We need to show that f is strictly convex. Since f is an infinitely differentiable function, we need to show that $f''(u) > 0$ for all $u \in \mathbb{R}$. First, note that the reflection formula (2.5) implies that

$$(2.10) \quad f(-u) = \frac{u}{2} + f(u).$$

Therefore

$$f''(-u) = f''(u).$$

As a result, we only need to show that $f''(u) > 0$ for all $u \in \mathbb{R}^+$. By definition,

$$g'(u) = \frac{\vartheta'(u)}{\vartheta(u)}, \quad g''(u) = \frac{\vartheta''(u)\vartheta(u) - \vartheta'(u)^2}{\vartheta(u)^2},$$

and

$$\begin{aligned} f''(u) &= e^{2u} g''(u) + e^u g'(e^u) \\ &= e^{2u} \frac{\vartheta''(e^u)\vartheta(e^u) - \vartheta'(e^u)^2 + e^{-u}\vartheta(e^u)\vartheta'(e^u)}{\vartheta(e^u)^2}. \end{aligned}$$

Hence, we have to prove that for all $v \geq 1$,

$$h(v) = \vartheta''(v)\vartheta(v) - \vartheta'(v)^2 + v^{-1}\vartheta(v)\vartheta'(v) > 0.$$

Again by definition,

$$(2.11)$$

$$\begin{aligned} h(v) &= \sum_{j \in \mathbb{Z}} \pi^2 j^4 e^{-\pi v j^2} \sum_{k \in \mathbb{Z}} e^{-\pi v k^2} - \sum_{j \in \mathbb{Z}} \pi j^2 e^{-\pi v j^2} \sum_{k \in \mathbb{Z}} \pi k^2 e^{-\pi v k^2} - v^{-1} \sum_{j \in \mathbb{Z}} e^{-\pi v j^2} \sum_{k \in \mathbb{Z}} \pi k^2 e^{-\pi v k^2} \\ &= \frac{\pi}{2} \sum_{(j,k) \in \mathbb{Z}^2} \left[\pi j^4 + \pi k^4 - 2\pi j^2 k^2 - \frac{1}{v}(j^2 + k^2) \right] e^{-\pi v(j^2 + k^2)}. \end{aligned}$$

For fixed $v \geq 1$ and $(j, k) \in \mathbb{Z}^2$, define

$$C(j, k) = \pi j^4 + \pi k^4 - 2\pi j^2 k^2 - \frac{1}{v}(j^2 + k^2) = \pi(k^2 - j^2)^2 - \frac{1}{v}(j^2 + k^2).$$

If $C(j, k) > 0$ for all $(j, k) \in \mathbb{Z}^2$, it follows directly that $h(v) > 0$. However, $C(k, k) = -2v^{-1}k^2 \leq 0$, which renders the verification of $h(v) > 0$ slightly more complicated. We need to compare the values of $C(j, k)$ for different pairs of (j, k) . Since $C(j, k) = C(-j, k) = C(j, -k) = C(-j, -k)$ and $C(j, k) = C(k, j)$, we need only to consider $C(j, k)$ as a function of $(j, k) \in \mathbb{N}_0$ with $j \leq k$. Here $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. We claim that

(a) If $j \neq k$, $C(j, k) > 0$,

(b) For any $k \geq 1$, $C(k-1, k) + \frac{1}{2}C(k, k) > 0$.

By rewriting $C(j, k)$ as

$$C(j, k) = \pi j^4 - \left(2\pi k^2 + \frac{1}{v} \right) j^2 + \pi k^4 - \frac{k^2}{v},$$

we see that for fixed k , $C(j, k)$ is decreasing if $j \in [0, k]$. Therefore, to prove (a), it suffices to verify that $C(k-1, k) > 0$. Since $C(k, k) \leq 0$, this will follow if we prove (b). Now for $v \geq 1$, the function

$$\begin{aligned} q(k) &= C(k-1, k) + \frac{1}{2}C(k, k) = \pi(4k^2 - 4k + 1) - \frac{1}{v}(2k^2 - 2k + 1) - \frac{k^2}{v} \\ &\geq \pi(4k^2 - 4k + 1) - (3k^2 - 2k + 1) \\ &= (4\pi - 3)k^2 - 2(2\pi - 1)k + \pi - 1 \end{aligned}$$

is increasing for $k \geq 1$, and $q(1) = \pi - 2 > 0$, this proves (b). Returning to the proof of $h(v) > 0$, we write the summation over $(j, k) \in \mathbb{Z}^2$ in (2.11) as

$$\frac{\pi}{2} \left\{ \sum_{\substack{(j,k) \in \mathbb{Z}^2 \\ |j-k| \geq 2}} + \sum_{\substack{(j,k) \in \mathbb{Z}^2 \\ |j-k|=1}} + \sum_{j=k} \right\} C(j, k) e^{-\pi v(j^2 + k^2)}.$$

By (a), the first sum is strictly positive. Now since $C(0, 0) = 0$ and $C(\pm j, \pm k) = C(j, k) = C(k, j)$, the last two sums can be written as

$$2\pi \sum_{k=1}^{\infty} C(k-1, k) e^{-\pi((k-1)^2 + k^2)} + \pi \sum_{k=1}^{\infty} C(k, k) e^{-2\pi k^2}.$$

Using the fact that e^{-x} is a decreasing function, we find that this term is larger than

$$2\pi \sum_{k=1}^{\infty} \left(C(k-1, 1) + \frac{1}{2}C(k, k) \right) e^{-2\pi k^2},$$

which, by (b), is positive. This concludes the proof that $h(v) > 0$, and therefore the assertion of the proposition follows. $\square$

3. THE SIGN OF EPSTEIN ZETA FUNCTION

In this section, we are going to study the sign of the Xi-function $\Xi_n(s; a_1, \dots, a_n)$ when $s \in \mathbb{R} \setminus \{0, n/2\}$. First, since $\pi^{-s}\Gamma(s) > 0$ for $s > 0$ and it is obvious from the power series definition of $Z_n(s; a_1, \dots, a_n)$ (1.4) that $Z_n(s; a_1, \dots, a_n) > 0$ for all $s > n/2$, we immediately deduce that

$$\Xi_n(s; a_1, \dots, a_n) > 0 \quad \text{for all } s > \frac{n}{2} \text{ and } (a_1, \dots, a_n) \in (\mathbb{R}^+)^n.$$

From the functional equation (1.5), we then obtain

$$\Xi_n(s; a_1, \dots, a_n) > 0 \quad \text{for all } s < 0 \text{ and } (a_1, \dots, a_n) \in (\mathbb{R}^+)^n.$$

In other words, $\Xi_n(s; a_1, \dots, a_n) > 0$ for all $s \in (-\infty, 0) \cup (n/2, \infty)$. For the remaining case where $s \in (0, n/2)$, the sign of the function $\Xi_n(s; a_1, \dots, a_n)$ is not trivial. Using the formula (2.1), we find that around the two simple poles $s = 0$ and $s = \frac{n}{2}$,

$$\begin{aligned} \Xi_n(s; a_1, \dots, a_n) &= -\frac{V}{s} + O(1) \quad \text{as } s \rightarrow 0, \\ \Xi_n(s; a_1, \dots, a_n) &= -\frac{V^{-1}}{\frac{n}{2} - s} + O(1) \quad \text{as } s \rightarrow \frac{n}{2}. \end{aligned}$$

Therefore we can conclude that

Proposition 3.1. *For any $a = (a_1, \dots, a_n) \in (\mathbb{R}^+)^n$, one can find a right nonempty neighbourhood $I_1(a) = (0, s_1(a))$ of $s = 0$ and a left nonempty neighbourhood $I_2(a) = (s_2(a), \frac{n}{2})$ of $n/2$ such that*

$$\Xi_n(s; a_1, \dots, a_n) < 0, \quad \text{for all } s \in I_1(a) \cup I_2(a).$$

In fact, when $n = 1$, using the fact that

$$\Xi_1(s; a) = \sqrt{a} \pi^{-s} \Gamma(s) Z_1(s; a) = 2a^{-2s+\frac{1}{2}} \pi^{-s} \Gamma(s) \zeta_R(2s),$$

it is easy to verify a stronger result:

Proposition 3.2.

$$\Xi_1(s; a) < 0, \quad \text{for all } s \in \left(0, \frac{1}{2}\right).$$

Proof. It suffices to show that $\zeta_R(s) < 0$ for all $s \in (0, 1)$. For $s > 0$, there is a well-known formula

$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k^s} = (1 - 2^{1-s}) \zeta_R(s).$$

The alternating series on the left hand side is convergent and positive for all $s > 0$. However, for $s \in (0, 1)$, $2^{1-s} > 1$. Therefore, $\zeta_R(s) < 0$ for all $s \in (0, 1)$. This completes the proof. $\square$

In the study of the zeros of Riemann zeta function, this result asserts that $\zeta_R(s)$ has no nontrivial zeros in $(0, 1)$, a necessary condition for the validity of Riemann hypothesis. For $n \geq 2$, in view of Proposition 3.1, we find that a necessary condition for the validity of the generalized Riemann hypothesis for $Z_n(s; a_1, \dots, a_n)$ at a fixed $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$, is that $\Xi_n(s; a_1, \dots, a_n) < 0$ for all $s \in (0, n/2)$. However, this is not true for all $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$. In fact, we can show that

Proposition 3.3. *Given $n \geq 2$, $j \in \{1, \dots, n\}$ and $s \in (0, n/2)$, if we fix $a_1, \dots, a_{j-1}, a_{j+1}, \dots, a_n$ and vary a_j , then as a_j is large enough, $\Xi_n(s; a_1, \dots, a_n)$ is positive.*

Proof. Since $\Xi_n(s; a_1, \dots, a_n)$ is symmetric in the variables $a_1, \dots, a_n$, it is sufficient to show that given $n \geq 2$ and $s \in (0, n/2)$, there exists $j \in \{1, \dots, n\}$ such that if we fix $a_1, \dots, a_{j-1}, a_{j+1}, \dots, a_n$ and vary a_j , then as a_j is large enough, $\Xi_n(s; a_1, \dots, a_n)$ is positive.

First we consider the case $s = \frac{m}{2}$, where $m \in \{1, 2, \dots, n-1\}$. Equation (2.2) and the fact that (see [18])

$$\begin{aligned} \pi^{-s} \Gamma(s) \zeta_R(2s) &= -\frac{1}{2} \left\{ \frac{1}{s} + \log(4\pi) + \psi(1) \right\} + O(s), & \text{as } s \rightarrow 0, \\ \pi^{-s} \Gamma(s) \zeta_R(2s) &= \frac{1}{2} \left\{ \frac{1}{s - \frac{1}{2}} - \log(4\pi) - \psi(1) \right\} + O\left(s - \frac{1}{2}\right), & \text{as } s \rightarrow \frac{1}{2}, \end{aligned}$$

give

$$\begin{aligned}
(3.1) \quad V^{-1}\Xi_n\left(\frac{m}{2}; a_1, \dots, a_n\right) &= \frac{1}{\prod_{l=1}^m a_l} \left\{ \log \frac{a_{m+1}}{a_m} - \log(4\pi) - \psi(1) \right\} \\
&+ 2 \sum_{j \in \{0, 1, \dots, n-1\} \setminus \{m-1, m\}} \frac{\pi^{\frac{j-m}{2}} \Gamma\left(\frac{m-j}{2}\right)}{a_{j+1}^{m-j} \prod_{l=1}^j a_l} \zeta_R(m-j) + 4 \sum_{j=1}^{n-1} \frac{1}{\prod_{l=1}^j a_l} \times \\
&\sum_{\mathbf{k} \in \mathbb{Z}^j \setminus \{\mathbf{0}\}} \sum_{p=1}^{\infty} \frac{1}{(pa_{j+1})^{\frac{m-j}{2}}} \left(\sum_{l=1}^j \left[\frac{k_l}{a_l} \right]^2 \right)^{\frac{m-j}{4}} K_{\frac{m-j}{2}} \left(2\pi pa_{j+1} \sqrt{\sum_{l=1}^j \left[\frac{k_l}{a_l} \right]^2} \right).
\end{aligned}$$

Since the function $\pi^{-s}\Gamma(s)\zeta_R(2s)$ is strictly positive for $s < 0$ or $s > \frac{1}{2}$ and the function $K_s(z)$ is positive for all $s \in \mathbb{R}$ and $z \in \mathbb{R}^+$, the last two terms of (3.1) are always positive. The sign of the first term depend on the ratio a_{m+1}/a_m . It is easy to see that if $a_1, \dots, a_m, a_{m+2}, \dots, a_n$ are kept fixed, then for a_{m+1} large, the first term is positive and therefore $\Xi_n\left(\frac{m}{2}; a_1, \dots, a_n\right) > 0$. This establishes the proposition when $s \in \{m/2 : m = 1, 2, \dots, n-1\}$.

For the general case, given $s \in (0, n/2) \setminus \{m/2 : m = 1, 2, \dots, n-1\}$, there is a unique $m \in \{0, \dots, n-1\}$ such that $0 < 2s - m < 1$. Equation (2.2) gives

$$\begin{aligned}
V^{-1}\Xi_n(s; a_1, \dots, a_n) &= \frac{\pi^{\frac{m}{2}-s}\Gamma\left(s - \frac{m}{2}\right)}{a_{m+1}^{2s-m} \prod_{l=1}^m a_l} \zeta_R(2s - m) \\
&+ 2 \sum_{j=0}^{m-1} \frac{\pi^{\frac{j}{2}-s}\Gamma\left(s - \frac{j}{2}\right)}{a_{j+1}^{2s-j} \prod_{l=1}^j a_l} \zeta_R(2s - j) + 2 \sum_{j=m+1}^{n-1} \frac{\pi^{\frac{j}{2}-s}\Gamma\left(s - \frac{j}{2}\right)}{a_{j+1}^{2s-j} \prod_{l=1}^j a_l} \zeta_R(2s - j) + 4 \sum_{j=1}^{n-1} \frac{1}{\prod_{l=1}^j a_l} \times \\
&\sum_{\mathbf{k} \in \mathbb{Z}^j \setminus \{\mathbf{0}\}} \sum_{p=1}^{\infty} \frac{1}{(pa_{j+1})^{s-\frac{j}{2}}} \left(\sum_{l=1}^j \left[\frac{k_l}{a_l} \right]^2 \right)^{\frac{s-j}{2}} K_{s-\frac{j}{2}} \left(2\pi pa_{j+1} \sqrt{\sum_{l=1}^j \left[\frac{k_l}{a_l} \right]^2} \right).
\end{aligned}$$

Only the first term is negative. If $a_1, \dots, a_m, a_{m+2}, \dots, a_n$ are fixed and $a_{m+1} \rightarrow \infty$, the second term is fixed but the first term approaches zero. Therefore, $\Xi_n(s; a_1, \dots, a_n) > 0$ when a_{m+1} is large enough, and our assertion is proved.

We would like to remark that the case $s = \frac{m}{2}$, where $m \in \{1, 2, \dots, n-1\}$ can actually be deduced from the general case $s \in (0, n/2) \setminus \{m/2 : m = 1, 2, \dots, n-1\}$ using the joint continuity of $\Xi_n(s; a_1, \dots, a_n)$ as a function of s and $(a_1, \dots, a_n)$. $\square$

This proposition shows that if $s \in (0, n/2)$, then when one of the ratios between the a_i 's, $1 \leq i \leq n$ is large, the Epstein zeta function $Z_n(s; a_1, \dots, a_n)$ is positive, and therefore the generalized Riemann hypothesis does not hold for this zeta function. It is then interesting to ask whether the Riemann hypothesis will still be valid for some $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$. Another interesting question which is related to our problem [18] of determining whether there exists symmetry breaking mechanisms in an interacting fractional Klein–Gordon field is, for what values of s , the Xi-function $\Xi_n(s; a_1, \dots, a_n)$ will be negative for some $(a_1, \dots, a_n)$. Recall that we have proved in section 2 that for any fixed s , the minimum of $Z_n(s; a_1, \dots, a_n)$, as a function of $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$ with $\prod_{i=1}^n a_i$ fixed, appears at the point $a_1 = \dots = a_n$. Therefore to answer these two questions, we need to study the function $\Xi_n(s) = \Xi_n(s; 1, \dots, 1)$ first.

From (2.1), we find that

$$(3.2) \quad \Xi_n(s) = -\frac{1}{s} - \frac{1}{\frac{n}{2} - s} + \int_1^\infty t^{s-1} (\vartheta(t)^n - 1) dt + \int_1^\infty t^{\frac{n}{2}-1-s} (\vartheta(t)^n - 1) dt.$$

In order to have a more unified treatment for all n , we define $\hat{\Xi}_n(s) = \Xi_n\left(\frac{ns}{2}\right)$. Then

$$(3.3) \quad \hat{\Xi}_n(s) = -\frac{2}{n} \left(\frac{1}{s} + \frac{1}{1-s} \right) + \int_1^\infty t^{\frac{ns}{2}-1} (\vartheta(t)^n - 1) dt + \int_1^\infty t^{\frac{n}{2}(1-s)-1} (\vartheta(t)^n - 1) dt.$$

This formula shows that for all $s \in (0, 1)$ and $n \in \mathbb{N}$,

$$(3.4) \quad \hat{\Xi}_{n+1}(s) > \hat{\Xi}_n(s).$$

Moreover, for any fixed s , since

$$\frac{2}{n} \left(\frac{1}{s} + \frac{1}{1-s} \right) \longrightarrow 0 \quad \text{as } n \rightarrow \infty,$$

and

$$\begin{aligned} & \int_1^\infty t^{\frac{ns}{2}-1} (\vartheta(t)^n - 1) dt + \int_1^\infty t^{\frac{n}{2}(1-s)-1} (\vartheta(t)^n - 1) dt \\ & \geq \int_1^\infty t^{\frac{s}{2}-1} (\vartheta(t)^n - 1) dt + \int_1^\infty t^{\frac{1}{2}(1-s)-1} (\vartheta(t)^n - 1) dt > 0, \end{aligned}$$

therefore $\hat{\Xi}_n(s)$ becomes positive when n is large enough. In fact for $s = 1/2$, a lengthy analysis (31 pages) has been done in [46] and it was proved that

Lemma 3.4.

$$\begin{cases} \Xi_n\left(\frac{n}{4}\right) = \hat{\Xi}_n\left(\frac{1}{2}\right) < 0, & \text{if } 1 \leq n \leq 9; \\ \Xi_n\left(\frac{n}{4}\right) = \hat{\Xi}_n\left(\frac{1}{2}\right) > 0, & \text{if } n \geq 10. \end{cases}$$

Proof. Here we provide a much shorter proof of this result. Using the inequality (3.4), we need only to show that

$$(3.5) \quad \Xi_9\left(\frac{9}{4}\right) < 0 \quad \text{and} \quad \Xi_{10}\left(\frac{5}{2}\right) > 0.$$

Using the incomplete gamma function

$$\Gamma(\beta, x) = \int_x^\infty t^{\beta-1} e^{-t} dt, \quad x > 0,$$

we can rewrite (3.2) as

$$(3.6) \quad \begin{aligned} \Xi_n(s) = & -\frac{1}{s} - \frac{1}{\frac{n}{2} - s} + \sum_{k \in \mathbb{Z}^n \setminus \{0\}} (\pi|k|^2)^{-s} \Gamma(s, \pi|k|^2) \\ & + \sum_{k \in \mathbb{Z}^n \setminus \{0\}} (\pi|k|^2)^{-\frac{n}{2}+s} \Gamma\left(\frac{n}{2} - s, \pi|k|^2\right), \end{aligned}$$

where for $k \in \mathbb{Z}^n$, $|k|^2 = \sum_{i=1}^n k_i^2$. We need to obtain upper and lower bounds for $\Gamma(\beta, x)$. Using integration by parts, it is easy to show that when $\beta > 0$,

$$\Gamma(\beta, x) = x^{\beta-1} e^{-x} \sum_{j=0}^m \frac{\prod_{i=1}^j (\beta - i)}{x^j} + \left[\prod_{i=1}^{m+1} (\beta - i) \right] \int_x^\infty t^{\beta-m-2} e^{-t} dt.$$

From this, we find that

$$(3.7) \quad \Gamma(\beta, x) \leq x^{\beta-1} e^{-x} \sum_{j=0}^{[\beta]} \frac{\prod_{i=1}^j (\beta - i)}{x^j}, \quad \Gamma(\beta, x) \geq x^{\beta-1} e^{-x} \sum_{j=0}^{[\beta]+1} \frac{\prod_{i=1}^j (\beta - i)}{x^j}.$$

Therefore from (3.6), we have

$$\Xi_9 \left(\frac{9}{4} \right) \leq -\frac{8}{9} + 2 \sum_{k \in \mathbb{Z}^9 \setminus \{0\}} (\pi |k|^2)^{-1} e^{-\pi |k|^2} \sum_{j=0}^2 \frac{\prod_{i=1}^j (\frac{9}{4} - i)}{(\pi |k|^2)^j}.$$

For the summation over $k \in \mathbb{Z}^9 \setminus \{0\}$, we single out the 18 terms that contribute to $|k| = 1$, i.e. we write

$$(3.8) \quad \begin{aligned} & \sum_{k \in \mathbb{Z}^9 \setminus \{0\}} (\pi |k|^2)^{-1} e^{-\pi |k|^2} \sum_{j=0}^2 \frac{\prod_{i=1}^j (\frac{9}{4} - i)}{(\pi |k|^2)^j} \\ &= \frac{18}{\pi} e^{-\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (\frac{9}{4} - i)}{\pi^j} + \sum_{k \in \mathbb{Z}^9 \setminus \{k : |k|=0 \text{ or } 1\}} (\pi |k|^2)^{-1} e^{-\pi |k|^2} \sum_{j=0}^2 \frac{\prod_{i=1}^j (\frac{9}{4} - i)}{(\pi |k|^2)^j}. \end{aligned}$$

The first term of (3.8) gives

$$(3.9) \quad \frac{18}{\pi} e^{-\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (\frac{9}{4} - i)}{\pi^j} = 0.3540;$$

whereas the second term of (3.8) is

$$(3.10) \quad \begin{aligned} & \sum_{k \in \mathbb{Z}^9 \setminus \{k : |k|=0 \text{ or } 1\}} (\pi |k|^2)^{-1} e^{-\pi |k|^2} \sum_{j=0}^2 \frac{\prod_{i=1}^j (\frac{9}{4} - i)}{(\pi |k|^2)^j} \\ & \leq \frac{1}{2\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (\frac{9}{4} - i)}{(2\pi)^j} \sum_{k \in \mathbb{Z}^9 \setminus \{k : |k|=0 \text{ or } 1\}} e^{-\pi |k|^2}. \end{aligned}$$

Using a simple inequality

$$\exp \left(-\pi \sum_{i=1}^n k_i^2 \right) \leq \exp \left(-\pi \sum_{i=1}^n |k_i| \right),$$

we find that

$$\begin{aligned} \sum_{k \in \mathbb{Z}^9 \setminus \{k : |k|=0 \text{ or } 1\}} e^{-\pi |k|^2} &= \sum_{k \in \mathbb{Z}^9} e^{-\pi |k|^2} - 1 - 18e^{-\pi} \\ &\leq \sum_{k \in \mathbb{Z}^9} e^{-\pi \sum_{i=1}^9 |k_i|} - 1 - 18e^{-\pi} \\ &= \left(1 + 2 \sum_{m=1}^{\infty} e^{-\pi m} \right)^9 - 1 - 18e^{-\pi} \\ &= \left(\frac{e^{\pi} + 1}{e^{\pi} - 1} \right)^9 - 1 - 18e^{-\pi}. \end{aligned}$$

Therefore, the term (3.10) is bounded above by

$$\frac{1}{2\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (\frac{9}{4} - i)}{(2\pi)^j} \left(\left(\frac{e^\pi + 1}{e^\pi - 1} \right)^9 - 1 - 18e^{-\pi} \right) = 0.0769.$$

Combine with (3.9), we find that (3.8) is bounded above by

$$0.3540 + 0.0769 = 0.4309 < \frac{4}{9}.$$

Therefore, $\Xi_9(9/4) < 0$. Now for $\Xi_{10}(5/2)$, a similar argument but using the lower bound in (3.7) for the incomplete gamma function gives

$$\begin{aligned} \Xi_{10} \left(\frac{5}{2} \right) &> -\frac{4}{5} + 2 \sum_{k \in \mathbb{Z}^{10} \setminus \{0\}} (\pi|k|^2)^{-1} e^{-\pi|k|^2} \sum_{j=0}^3 \frac{\prod_{i=1}^j (\frac{5}{2} - i)}{(\pi|k|^2)^j} \\ &> -\frac{4}{5} + \frac{40}{\pi} e^{-\pi} \sum_{j=0}^3 \frac{\prod_{i=1}^j (\frac{5}{2} - i)}{\pi^j} = 0.04808 > 0. \end{aligned}$$

As a side remark, using standard mathematics softwares such as MATLAB and MATHEMATICA, it is easy to compute $\Xi_n(s)$ to any desired degree of accuracy from formula (3.2) or (3.6). Up to 10^{-12} , we have

$$\begin{aligned} \Xi_9 \left(\frac{9}{4} \right) &= -0.065884758538, \\ \Xi_{10} \left(\frac{5}{2} \right) &= 0.205903040487. \end{aligned}$$

□

Lemma 3.4, Proposition 3.1 and the inequality (3.4) immediately implies that

Proposition 3.5. *For all $n \geq 10$, the sets*

$$(3.11) \quad I_{n,+} = \{s \in (0, n/2) : \Xi_n(s) > 0\},$$

$$(3.12) \quad I_{n,-} = \{s \in (0, n/2) : \Xi_n(s) < 0\}$$

are open nonempty subsets of $(0, n/2)$. Moreover, $n/4 \in I_{n,+}$.

In fact, from the functional equation $\Xi_n(s) = \Xi_n((n/2) - s)$, one can even conclude that the intervals $I_{n,+}$ and $I_{n,-}$ are symmetric with respect to the point $n/4$, i.e.

$$\frac{n}{2} - I_{n,+} := \left\{ \frac{n}{2} - s : s \in I_{n,+} \right\} = I_{n,+}$$

and similarly for $I_{n,-}$. Since

$$\lim_{s \rightarrow 0^+} \Xi_n(s) = -\infty,$$

there must exists an odd number of points $\gamma_{n,1}, \dots, \gamma_{n,2m_n+1}$ so that $0 < \gamma_{n,1} < \gamma_{n,2} \leq \gamma_{n,3} < \dots < \gamma_{n,2m_n} \leq \gamma_{n,2m_n+1} < n/4$ and

$$I_{n,+} = \bigcup_{i=1}^{m_n} (\gamma_{n,2i-1}, \gamma_{n,2i}) \bigcup \left(\gamma_{n,2m_n+1}, \frac{n}{2} - \gamma_{n,2m_n+1} \right) \bigcup_{i=1}^{m_n} \left(\frac{n}{2} - \gamma_{n,2i}, \frac{n}{2} - \gamma_{n,2i-1} \right).$$

For $n \leq 9$, Lemma 3.4 is not sufficient to show that $\Xi_n(s) < 0$ for all $s \in (0, n/2)$. Nevertheless, one can employ the same method as the proof in Lemma 3.4 to show that

Proposition 3.6. $\Xi_9(s) < 0$ for all $s \in (0, 9/2)$.

Then the inequality (3.4) shows that

Proposition 3.7. For all $1 \leq n \leq 9$, the function $\Xi_n(s)$ is negative for all $s \in (0, n/2)$.

We give the proof of Proposition 3.6 here.

Proof. By the reflection formula $\Xi_9(s) = \Xi_9((9/2) - s)$, it is sufficient to show that $\Xi_9(s) < 0$ for all $s \in (0, 9/4]$. Let

$$\mathcal{Z}_1(s) = -\frac{1}{s} - \frac{1}{\frac{9}{2} - s}$$

and

$$\mathcal{Z}_2(s) = \int_1^\infty t^{s-1} (\vartheta(t)^9 - 1) dt + \int_1^\infty t^{\frac{9}{2}-s-1} (\vartheta(t)^9 - 1) dt,$$

so that $\Xi_9(s) = \mathcal{Z}_1(s) + \mathcal{Z}_2(s)$. By taking derivatives with respect to s , it is easy to verify that as functions of s , $\mathcal{Z}_1(s)$ is increasing on $(0, 9/4]$ and $\mathcal{Z}_2(s)$ is decreasing on $[0, 9/4]$. Using the arguments in the proof of Lemma 3.4, we find that

$$\begin{aligned} \mathcal{Z}_2(0) &\leq \frac{18}{\pi} e^{-\pi} + \frac{1}{2\pi} \left(\left(\frac{e^\pi + 1}{e^\pi - 1} \right)^9 - 1 - 18e^{-\pi} \right) \\ &\quad + \frac{18}{\pi} e^{-\pi} \sum_{j=0}^4 \frac{\prod_{i=1}^j (\frac{9}{2} - i)}{\pi^j} + \frac{1}{2\pi} \sum_{j=0}^4 \frac{\prod_{i=1}^j (\frac{9}{2} - i)}{(2\pi)^j} \left(\left(\frac{e^\pi + 1}{e^\pi - 1} \right)^9 - 1 - 18e^{-\pi} \right) \\ &= 1.2926. \end{aligned}$$

Now we want to find an x so that $\mathcal{Z}_1(x) < -1.2926$. By try and error method, we find that $x = 0.95$ satisfies the required condition. In fact,

$$\mathcal{Z}_1(0.95) = -1.3343.$$

Therefore, for all $s \in (0, 0.95]$,

$$\Xi_9(s) = \mathcal{Z}_1(s) + \mathcal{Z}_2(s) \leq \mathcal{Z}_1(0.95) + \mathcal{Z}_2(0) \leq -0.0417 < 0.$$

Repeating these steps, we have first

$$\begin{aligned} \mathcal{Z}_2(0.95) &\leq \frac{18}{\pi} e^{-\pi} + \frac{1}{2\pi} \left(\left(\frac{e^\pi + 1}{e^\pi - 1} \right)^9 - 1 - 18e^{-\pi} \right) \\ &\quad + \frac{18}{\pi} e^{-\pi} \sum_{j=0}^3 \frac{\prod_{i=1}^j (3.55 - i)}{\pi^j} + \frac{1}{2\pi} \sum_{j=0}^3 \frac{\prod_{i=1}^j (3.55 - i)}{(2\pi)^j} \left(\left(\frac{e^\pi + 1}{e^\pi - 1} \right)^9 - 1 - 18e^{-\pi} \right) \\ &= 0.9728. \end{aligned}$$

Then by try and error, we find that

$$\mathcal{Z}_1(1.55) = -0.9841.$$

Therefore, for all $s \in [0.95, 1.55]$,

$$\Xi_9(s) = \mathcal{Z}_1(s) + \mathcal{Z}_2(s) \leq \mathcal{Z}_1(1.55) + \mathcal{Z}_2(0.95) \leq -0.0113 < 0.$$




FIGURE 1. The graphs of $\hat{\Xi}_n(s) = \Xi_n\left(\frac{ns}{2}\right)$ for $s \in (0, 1)$.

Repeating again, we find that

$$\begin{aligned} \mathcal{Z}_2(1.55) &\leq \frac{18}{\pi} e^{-\pi} \sum_{j=0}^1 \frac{\prod_{i=1}^j (1.55 - i)}{\pi^j} + \frac{1}{2\pi} \sum_{j=0}^1 \frac{\prod_{i=1}^j (1.55 - i)}{(2\pi)^j} \left(\left(\frac{e^\pi + 1}{e^\pi - 1} \right)^9 - 1 - 18e^{-\pi} \right) \\ &\quad + \frac{18}{\pi} e^{-\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (2.95 - i)}{\pi^j} + \frac{1}{2\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (2.95 - i)}{(2\pi)^j} \left(\left(\frac{e^\pi + 1}{e^\pi - 1} \right)^9 - 1 - 18e^{-\pi} \right) \\ &= 0.8943, \end{aligned}$$

and

$$\mathcal{Z}_1(2) = -0.9.$$

Therefore, for all $s \in [1.55, 2]$,

$$\Xi_9(s) = \mathcal{Z}_1(s) + \mathcal{Z}_2(s) \leq \mathcal{Z}_1(2) + \mathcal{Z}_2(1.55) \leq -0.0057 < 0.$$

Finally,

$$\begin{aligned} \mathcal{Z}_2(2) &\leq \frac{18}{\pi} e^{-\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (2 - i)}{\pi^j} + \frac{1}{2\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (2 - i)}{(2\pi)^j} \left(\left(\frac{e^\pi + 1}{e^\pi - 1} \right)^9 - 1 - 18e^{-\pi} \right) \\ &\quad + \frac{18}{\pi} e^{-\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (2.5 - i)}{\pi^j} + \frac{1}{2\pi} \sum_{j=0}^2 \frac{\prod_{i=1}^j (2.5 - i)}{(2\pi)^j} \left(\left(\frac{e^\pi + 1}{e^\pi - 1} \right)^9 - 1 - 18e^{-\pi} \right) \\ &= 0.8649, \end{aligned}$$

and

$$\mathcal{Z}_1(2.25) = -0.8889,$$

which implies that for all $s \in [2, 2.25]$,

$$\Xi_9(s) = \mathcal{Z}_1(s) + \mathcal{Z}_2(s) \leq \mathcal{Z}_1(2.25) + \mathcal{Z}_2(2) \leq -0.0240 < 0.$$

This completes the proof of the proposition. $\square$

In Figure 1, we show the graphs of $\hat{\Xi}_n(s) = \Xi_n(ns/2)$ for $s \in (0, 1)$ and $1 \leq n \leq 10$. From the graphs, we see that for $1 \leq n \leq 10$, $\hat{\Xi}_n(s)$ is increasing for $s \in (0, 1/2)$, decreasing for $s \in (1/2, 1)$, and $s = 1/2$ is a local maximum of $\hat{\Xi}_n(s)$. One would




FIGURE 2. The graphs of $\hat{\Xi}_{11}(s) = \Xi_{11}\left(\frac{11s}{2}\right)$ and $\hat{\Xi}_{12}(s) = \Xi_{12}(6s)$.

tend to jump into the conclusion that this will be true for all n . However, this is not the case. Since $\hat{\Xi}_n(s)$ satisfies the functional equation $\hat{\Xi}_n(s) = \hat{\Xi}_n(1-s)$, it is easy to verify that $\hat{\Xi}'_n(1/2) = 0$ and therefore $s = 1/2$ is indeed a local extremum of $\hat{\Xi}_n(s)$. In order to determine the nature of this extremum, it is necessary to look at the second derivative of $\hat{\Xi}_n(s)$. We obtain from (3.3) that

$$(3.13) \quad \begin{aligned} \hat{\Xi}''_n(s) = & -\frac{4}{n} \left(\frac{1}{s^3} + \frac{1}{(1-s)^3} \right) + \left(\frac{n}{2} \right)^2 \int_1^\infty t^{\frac{ns}{2}-1} (\log t)^2 (\vartheta(t)^n - 1) dt \\ & + \left(\frac{n}{2} \right)^2 \int_1^\infty t^{\frac{n}{2}(1-s)-1} (\vartheta(t)^n - 1) dt. \end{aligned}$$

From here, it is easy to verify that

$$(3.14) \quad \hat{\Xi}''_{n+1}(s) > \hat{\Xi}''_n(s).$$

Using the same argument as employed in showing that for any fixed $s \in (0, n/2)$, $\hat{\Xi}_n(s) > 0$ when n is large enough, we can use (3.13) to prove that for any fixed $s \in (0, n/2)$, $\Xi''_n(s) > 0$ whenever n is large enough. This implies that the point $n/4$ will become a local minimum of $\Xi_n(s)$ when n is large enough. In fact, using numerical computation, we find that

$$\begin{aligned} \Xi''_{10}\left(\frac{5}{2}\right) &= -0.101080515709, \\ \Xi''_{11}\left(\frac{11}{4}\right) &= 0.009568954836, \end{aligned}$$

which together with (3.14), imply that for $1 \leq n \leq 10$, $s = n/4$ is a local maximum of $\Xi_n(s)$; whereas for $n \geq 11$, $s = n/4$ is a local minimum of $\Xi_n(s)$. In Figure 2, we show graphically that $s = 1/2$ is a local minimum of $\hat{\Xi}_{11}(s)$ and $\hat{\Xi}_{12}(s)$.

In Table 1, we tabulate the interval $I_{n,+}$, $10 \leq n \leq 21$ where $\Xi_n(s)$ is positive.

Table 1: The interval $I_{n,+}$ for $10 \leq n \leq 21$.

n	$I_{n,+}$	n	$I_{n,+}$
10	(1.0899, 3.9101)	11	(0.6401, 4.8599)
12	(0.3976, 5.6024)	13	(0.2498, 6.2502)
14	(0.1562, 6.8438)	15	(0.0964, 7.4036)
16	(0.0585, 7.9415)	17	(0.0348, 8.4652)
18	(0.0202, 8.9798)	19	(0.0115, 9.4885)
20	(0.0064, 9.9936)	21	(0.0034, 10.4966)

One notices that for $10 \leq n \leq 21$, $m_n = 0$, $I_{n,+} = (\gamma_{n,1}, (n/2) - \gamma_{n,1})$, $\gamma_{n,1}$ is decreasing and $I_{n,+} \subseteq I_{n+1,+}$. In fact, from the formula (3.2), it is easy to verify that for $s \in (0, n/2)$,

$$\Xi_{n+1}(s) > \Xi_n(s).$$

This shows that $I_{n,+} \subseteq I_{n+1,+}$ for all $n \geq 10$. From this table, it is also natural to conjecture that $I_{n,+}$ is an open connected interval with center at $n/4$ for all $n \geq 10$. However, to prove this would require a very detailed analysis of $\Xi_n(s)$ and its higher derivatives. We do not intend to deal further into this problem here.

We now return to the discussion about the sign of the general Xi-function $\Xi_n(s; a_1, \dots, a_n)$ in the range $s \in (0, n/2)$. Since we have shown in Theorem 2.2 that for any fixed s , the point $a_1 = \dots = a_n$ is the minimum of $\Xi_n(s; a_1, \dots, a_n)$, $(a_1, \dots, a_n) \in (\mathbb{R}^n)^+$ with $\prod_{i=1}^n a_n$ fixed, now together with Proposition 3.5, Proposition 3.7 and Proposition 3.1, we obtain

Proposition 3.8. *For $1 \leq n \leq 9$, there exists a nonempty region containing the ray $a_1 = \dots = a_n$ in $(\mathbb{R}^n)^+$ where $\Xi_n(s; a_1, \dots, a_n)$ is negative for all $s \in (0, n/2)$.*

Proposition 3.9. *If $n \geq 10$, then for every $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$, there exists a nonempty open subset of $(0, n/2)$ where $\Xi_n(s; a_1, \dots, a_n)$ is positive and a nonempty open subset of $(0, n/2)$ where $\Xi_n(s; a_1, \dots, a_n)$ is negative.*

This proposition implies that generalized Riemann hypothesis is not true for all the Epstein zeta function of the form $Z_n(s; a_1, \dots, a_n)$ (i.e. Epstein zeta function of positive definite diagonal quadratic forms) when $n \geq 10$.

Now consider the Epstein zeta function as a function of $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$ with s fixed, we can conclude with the help of Propositions 3.3 and 3.7 that

Proposition 3.10. *If $1 \leq n \leq 9$, then for any fixed $s \in (0, n/2)$, there exists a nonempty open region $\Omega_{s,n}^+$ of $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$ where $\Xi_n(s; a_1, \dots, a_n)$ is positive and a nonempty open region $\Omega_{s,n}^-$ of $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$ where $\Xi_n(s; a_1, \dots, a_n)$ is negative. The ray $a_1 = \dots = a_n$ lies inside the region $\Omega_{s,n}^-$.*

Proposition 3.11. *If $n \geq 10$, then if $s \in I_{n,+}$ (eq. (3.11)), the function $\Xi_n(s; a_1, \dots, a_n)$ is positive for all $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$; whereas if $s \in I_{n,-}$ (eq. (3.12)), there exists a nonempty open region $\Omega_{s,n}^+$ of $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$ where $\Xi_n(s; a_1, \dots, a_n)$ is positive and a nonempty open region $\Omega_{s,n}^-$ of $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$ where $\Xi_n(s; a_1, \dots, a_n)$ is negative. In the latter case, the ray $a_1 = \dots = a_n$ lies inside the region $\Omega_{s,n}^-$.*

Propositions 3.10 and 3.11 are important for our study of symmetry breaking of interacting fractional Klein–Gordon field in [18]. In the following section, it will be shown that the region $\Omega_{s,n}^-$ is connected. Moreover, under the map $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n \mapsto (\log a_1, \dots, \log a_n) \in \mathbb{R}^n$, the projection of its image to the plane $x_1 + \dots + x_n = \text{constant}$ is a convex set.

4. CONVEXITY OF THE EPSTEIN ZETA FUNCTION

In this section, we show that

Theorem 4.1. *For any fixed s , when $\sum_{i=1}^n c_i = c$ is fixed, the function*

$$(4.1) \quad \begin{aligned} (c_1, \dots, c_n) &\mapsto \Xi_n(s; e^{c_1}, \dots, e^{c_n}) \\ &= -\frac{e^{\frac{s}{2}}}{s} - \frac{e^{-\frac{s}{2}}}{\frac{n}{2} - s} + e^{\frac{s}{2}} \int_1^\infty t^{s-1} \left(\prod_{i=1}^n \vartheta(e^{2c_i t}) - 1 \right) dt \\ &\quad + e^{-\frac{s}{2}} \int_1^\infty t^{\frac{n}{2}-s-1} \left(\prod_{i=1}^n \vartheta(e^{-2c_i t}) - 1 \right) dt, \end{aligned}$$

regarded as a function of any $(n-1)$ of the variables $\{c_1, \dots, c_n\}$, is a convex function.

Recall that a function $f(x)$, $x \in \mathbb{R}^n$ is convex if and only if for any $\lambda \in (0, 1)$ and any two points $x, y \in \mathbb{R}^n$,

$$f(\lambda x + (1-\lambda)y) \leq \lambda f(x) + (1-\lambda)f(y).$$

If f has continuous second partial derivatives, then f is convex if and only if the Hessian of f ,

$$H_f(x_1, \dots, x_n) = \left(\frac{\partial^2 f}{\partial x_i \partial x_j} \right)_{\substack{1 \leq i \leq n \\ 1 \leq j \leq n}},$$

is a positive semi-definite matrix. One way to prove that a Hermitian matrix is positive semi-definite is to use the Sylvester criterion, which says that an $n \times n$ Hermitian matrix is positive semi-definite if and only if all its leading principal minors are nonnegative. For $1 \leq i \leq n$, the i^{th} leading principal minor of an $n \times n$ matrix A is the determinant of the $i \times i$ square matrix on the upper left corner of A .

In the formula (4.1), the dependence on $(c_1, \dots, c_n)$ only comes from the terms $\prod_{i=1}^n \vartheta(e^{2c_i t})$ and $\prod_{i=1}^n \vartheta(e^{-2c_i t})$. Notice that if a function $f(x_1, \dots, x_n)$ is convex, then under any affine change of coordinates

$$x_i = \sum_{j=1}^n l_{ij} y_j + d_i, \quad 1 \leq i \leq n,$$

the function

$$\hat{f}(y_1, \dots, y_n) = f \left(\sum_{j=1}^n l_{1j} y_j + d_1, \dots, \sum_{j=1}^n l_{nj} y_j + d_n \right)$$

is also convex. Moreover, the projection of f to any hyperplane $\sum_{i=1}^n b_i x_i = b$ is also convex. Therefore, if the function

$$\Theta_n(x_1, \dots, x_n) = \prod_{i=1}^n \vartheta(e^{x_i}) = \prod_{i=1}^n \Theta_1(x_i), \quad (x_1, \dots, x_n) \in \mathbb{R}^n$$

is convex, then $\prod_{i=1}^n \vartheta(e^{2c_i t})$ and $\prod_{i=1}^n \vartheta(e^{-2c_i t})$, regarded as functions of $\{c_1, \dots, c_{n-1}\}$ (so that $c_n = c - \sum_{i=1}^{n-1} c_i$), are convex. This will show that $\Xi_n(s; e^{c_1}, \dots, e^{c_n})$ is a convex function of $(c_1, \dots, c_{n-1})$. Therefore to prove Theorem 4.1, we only need to

show that for all $n \geq 1$, the function $\Theta_n(x_1, \dots, x_n)$ is convex. For $n = 1$, we use the following fact:

Lemma 4.2. *If $g : \mathbb{R} \rightarrow \mathbb{R}$ is a convex function, then the function f defined by*

$$f(x) = e^{g(x)},$$

is also convex.

Proof. This is a simple consequence of the fact that the exponential function e^x is a convex function, and composition of convex functions are convex. $\square$

Using this lemma, we can immediately conclude from Proposition 2.1 that

Lemma 4.3. *The function*

$$\Theta_1(x) = \vartheta(e^x), \quad x \in \mathbb{R}$$

is convex.

For $n \geq 2$, in view of the definition of Θ_n , we are led to the study of a general function of the type

$$(4.2) \quad \mathcal{F}_n(x_1, \dots, x_n) = \prod_{i=1}^n f(x_i),$$

where $f : \mathbb{R} \rightarrow \mathbb{R}^+$ is a positive function of one variable. For such functions, its Hessian is given by

$$H_{\mathcal{F}_n}(x_1, \dots, x_n) = \mathcal{F}_n(x_1, \dots, x_n)^n \begin{pmatrix} \frac{f''(x_1)}{f(x_1)} & \frac{f'(x_1)f'(x_2)}{f(x_1)f(x_2)} & \cdots & \frac{f'(x_1)f'(x_n)}{f(x_1)f(x_n)} \\ \frac{f'(x_2)f'(x_1)}{f(x_2)f(x_1)} & \frac{f''(x_2)}{f(x_2)} & \cdots & \frac{f'(x_2)f'(x_n)}{f(x_2)f(x_n)} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{f'(x_n)f'(x_1)}{f(x_n)f(x_1)} & \frac{f'(x_n)f'(x_2)}{f(x_n)f(x_2)} & \cdots & \frac{f''(x_n)}{f(x_n)} \end{pmatrix}.$$

By the Sylvester criterion, to show that $\mathcal{F}_n$ is convex, it is sufficient to show that all the leading principal minors of

$$(4.3) \quad H_n(x_1, \dots, x_n) = \begin{pmatrix} \frac{f''(x_1)}{f(x_1)} & \frac{f'(x_1)f'(x_2)}{f(x_1)f(x_2)} & \cdots & \frac{f'(x_1)f'(x_n)}{f(x_1)f(x_n)} \\ \frac{f'(x_2)f'(x_1)}{f(x_2)f(x_1)} & \frac{f''(x_2)}{f(x_2)} & \cdots & \frac{f'(x_2)f'(x_n)}{f(x_2)f(x_n)} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{f'(x_n)f'(x_1)}{f(x_n)f(x_1)} & \frac{f'(x_n)f'(x_2)}{f(x_n)f(x_2)} & \cdots & \frac{f''(x_n)}{f(x_n)} \end{pmatrix}$$

are nonnegative. Notice that the i^{th} leading principal minor of $H_n(x_1, \dots, x_n)$ is the determinant of $H_i(x_1, \dots, x_i)$. Therefore, one only have to show that $\det H_n \geq 0$ for all $n \geq 1$. For $f(x) = \vartheta(e^x)$, the $n = 1$ case is already proved in Lemma 4.3. For $n \geq 2$, to further simplify our problem, notice that for fixed $(x_1, \dots, x_n) \in \mathbb{R}^n$, the matrix H_n has the form

$$J_n = \begin{pmatrix} z_1 & w_1 w_2 & \cdots & w_1 w_n \\ w_2 w_1 & z_2 & \cdots & w_2 w_n \\ \vdots & \vdots & \ddots & \vdots \\ w_n w_1 & w_n w_2 & \cdots & z_n \end{pmatrix}$$

for some $2n$ real numbers $z_1, \dots, z_n$ and $w_1, \dots, w_n$. When $n = 2$, the determinant of J_2 is

$$(4.4) \quad \det J_2 = z_1 z_2 - w_1^2 w_2^2 = (z_1 - w_1^2)(z_2 - w_2^2) + w_1^2(z_1 - w_1^2) + w_2^2(z_2 - w_2^2).$$

The reason for expressing the determinant in this form is inspired by the fact that for $z_i = \frac{f''(x_i)}{f(x_i)}$, $w_i = \frac{f'(x_i)}{f(x_i)}$,

$$(4.5) \quad z_i - w_i^2 = \frac{d^2}{dx_i^2} \log f(x_i).$$

Eq. (4.4) implies that if $z_i - w_i^2 \geq 0$, then $\det J_2 \geq 0$. This gives

Lemma 4.4. *Let $f : \mathbb{R} \rightarrow \mathbb{R}^+$ be a twice continuously differentiable positive function. If $\log f(x)$ is a convex function, then the function $\mathcal{F}_2(x_1, x_2) = f(x_1)f(x_2)$ is convex.*

Proof. Recall that $\mathcal{F}_2(x_1, x_2)$ is convex if and only if

$$\det H_1(x_1) = \frac{f''(x_1)}{f(x_1)} \geq 0$$

and

$$\det H_2(x_1, x_2) = \det J_2 \left[z_1 = \frac{f''(x_1)}{f(x_1)}, z_2 = \frac{f''(x_2)}{f(x_2)}; w_1 = \frac{f'(x_1)}{f(x_1)}, w_2 = \frac{f'(x_2)}{f(x_2)} \right] \geq 0.$$

As discussed above, if $\log f(x)$ is convex, then $\det H_2(x_1, x_2) \geq 0$. On the other hand, Lemma 4.2 implies that $f(x)$ is also convex. Therefore, $\det H_1(x_1) \geq 0$. This completes the proof. $\square$

For the case under consideration, $f(x) = \vartheta(e^x)$, and it is shown in Proposition 2.1 that $\log \vartheta(e^x)$ is a convex function. Therefore, Lemma 4.4 implies that

Lemma 4.5. *The function*

$$\Theta_2(x_1, x_2) = \vartheta(e^{x_1})\vartheta(e^{x_2}), \quad (x_1, x_2) \in \mathbb{R}^2$$

is convex.

Returning to the general case. First we claim the following, which is a generalization of (4.4).

Proposition 4.6. *For $n \geq 2$,*

$$(4.6) \quad \det J_n = \det \begin{pmatrix} z_1 & w_1 w_2 & \dots & w_1 w_n \\ w_2 w_1 & z_2 & \dots & w_2 w_n \\ \vdots & \vdots & \ddots & \vdots \\ w_n w_1 & w_n w_2 & \dots & z_n \end{pmatrix} = \prod_{i=1}^n (z_i - w_i^2) + \sum_{i=1}^n w_i^2 \left[\prod_{j \neq i} (z_j - w_j^2) \right].$$

Therefore, if $z_i \geq w_i^2$ for all $1 \leq i \leq n$, then $\det J_n \geq 0$.

We would like to discuss the consequence of this proposition first, and give the proof later. From the above discussion, the function $\mathcal{F}_n$ of the form (4.2) is convex if and only if all the determinants of H_i (eq. (4.3)), $1 \leq i \leq n$ are nonnegative. Now H_n is a matrix of the form J_n , with $z_i - w_i^2$ given by (4.5). Therefore, we conclude from Proposition 4.6 that

Proposition 4.7. *Let $f : \mathbb{R} \rightarrow \mathbb{R}^+$ be a twice continuously differentiable positive function. If $\log f(x)$ is a convex function, then the function*

$$\mathcal{F}_n(x_1, \dots, x_n) = \prod_{i=1}^n f(x_i)$$

is convex.

As in Lemma 4.5, this proposition implies that

Proposition 4.8. *For any $n \geq 1$, the function*

$$\Theta_n(x_1, \dots, x_n) = \prod_{i=1}^n \vartheta(e^{x_i}), \quad (x_1, \dots, x_n) \in \mathbb{R}^n,$$

is a convex function.

As discussed in the beginning of this section, this concludes the proof of Theorem 4.1. Therefore, what is left is the proof of Proposition 4.6.

Proof. By factoring out w_i from row i and column i , we have

$$\det J_n = \left[\prod_{i=1}^n w_i^2 \right]^2 \mathcal{J}_n \left(\frac{z_1}{w_1^2}, \frac{z_2}{w_2^2}, \dots, \frac{z_n}{w_n^2} \right),$$

where

$$\mathcal{J}_n(\alpha_1, \dots, \alpha_n) = \det \begin{pmatrix} \alpha_1 & 1 & 1 & \dots & 1 \\ 1 & \alpha_2 & 1 & \dots & 1 \\ 1 & 1 & \alpha_3 & \dots & 1 \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & 1 & 1 & \dots & \alpha_n \end{pmatrix}.$$

By elementary row operation, we find that

$$\begin{aligned} \mathcal{J}_n(\alpha_1, \dots, \alpha_n) &= \det \begin{pmatrix} \alpha_1 & 1 & 1 & \dots & 1 \\ 1 & \alpha_2 & 1 & \dots & 1 \\ 1 & 1 & \alpha_3 & \dots & 1 \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & 1 & 1 & \dots & \alpha_n \end{pmatrix} \\ &= \det \begin{pmatrix} \alpha_1 - 1 & 1 - \alpha_2 & 0 & \dots & 0 \\ 0 & \alpha_2 - 1 & 1 - \alpha_3 & \dots & 0 \\ 0 & 0 & \alpha_3 - 1 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & 1 & 1 & \dots & \alpha_n \end{pmatrix} \\ &= (\alpha_1 - 1) \mathcal{J}_{n-1}(\alpha_2, \dots, \alpha_n) + (-1)^{n-1} \prod_{i=2}^n (1 - \alpha_i). \end{aligned}$$

It then follows by induction that

$$\mathcal{J}_n(\alpha_1, \dots, \alpha_n) = \prod_{i=1}^n (\alpha_i - 1) + \sum_{i=1}^n \prod_{j \neq i} (\alpha_j - 1).$$

This proves the proposition. $\square$

Next we discuss the consequences of Theorem 4.1. Let j be a positive integer less than or equal to $n - 1$, A an $n \times j$ matrix such that the sum of every column vanishes, i.e.,

$$(4.7) \quad \sum_{i=1}^n A_{ij} = 0,$$

and let v be a vector in $\mathbb{R}^n$. Define a j -variable function $\Xi_{n,A,v}(s; b_1, \dots, b_j)$ by

$$\Xi_{n,A,v}(s; b_1, \dots, b_j) = \Xi_n \left(s; \exp \left[v_1 + \sum_{l=1}^j A_{1l} b_l \right], \dots, \exp \left[v_n + \sum_{l=1}^j A_{nl} b_l \right] \right).$$

Notice that the condition (4.7) implies

$$\prod_{i=1}^n \exp \left[v_i + \sum_{l=1}^j A_{il} b_l \right] = \exp \left[\sum_{i=1}^n v_i \right] = \text{constant}.$$

Since affine change of coordinates does not affect the convexity of a function, we conclude from Theorem 4.1 that $\Xi_{n,A,v}(s; b_1, \dots, b_j)$ is a convex function of $(b_1, \dots, b_j)$. As an example, if $v = 0$ and A is the $n \times (n - 1)$ matrix

$$(4.8) \quad A = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 1 \\ -1 & -1 & \dots & -1 \end{pmatrix},$$

then $\Xi_{n,A,v}(b_1, \dots, b_n)$ is just the function defined in (4.1) with $b_i = c_i$ for $1 \leq i \leq n - 1$ and $\sum_{i=1}^n c_n = 0$.

By the convexity of $\Xi_{n,A,v}(s; b_1, \dots, b_j)$, we can conclude that the region $\Omega_{s,A,v}^-$ of $(b_1, \dots, b_j) \in \mathbb{R}^j$ where $\Xi_{n,A,v}(s; b_1, \dots, b_j) < 0$ is a convex, and therefore connected region. Using the fact that at fixed $\prod_{i=1}^n a_i$, the minimum of $\Xi_n(s; a_1, \dots, a_n)$ appears at $a_1 = \dots = a_n$, one can even conclude that if $\Omega_{s,A,v}^-$ is nonempty and if $b = \hat{b}$ is a solution of the system

$$(4.9) \quad Ab + v = \left[\frac{1}{n} \sum_{i=1}^n v_i \right] \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix},$$

then $\hat{b} \in \Omega_{s,A,v}^-$. When A is given by (4.8) and $v = 0$, the system (4.9) has a unique solution $b = 0$. Therefore, the region of $(c_1, \dots, c_{n-1}) \in \mathbb{R}^n$ where the function $\Xi_n(s; e^{c_1}, \dots, e^{c_{n-1}}, e^{c - c_1 - \dots - c_{n-1}})$ (4.1) is negative is a convex connected region containing the origin $c_1 = \dots = c_{n-1} = 0$.

As a second example, suppose the variables $(a_1, \dots, a_n)$ in $\Xi_n(s; a_1, \dots, a_n)$ are such that $a_1 : \dots : a_n = 1 : k_2 : \dots : k_n$ and $\prod_{i=1}^n a_i = 1$. A simple computation shows that under these conditions, $a_1, \dots, a_n$ can be expressed as functions of $(k_2, \dots, k_n)$:

$$a_1 = \frac{1}{\prod_{i=2}^n k_i^{\frac{1}{k_i}}}, \quad a_2 = \frac{k_2^{\frac{n-1}{k_2}}}{\prod_{i=3}^n k_i^{\frac{1}{k_i}}}, \quad \dots, \quad a_n = \frac{k_n^{\frac{n-1}{k_n}}}{\prod_{i=2}^{n-1} k_i^{\frac{1}{k_i}}}.$$

Therefore the function $\Xi_n(s; a_1, \dots, a_n)$ can be regarded as a function of $\log k_i$, $2 \leq i \leq n$ with corresponding A and v given by

$$A = \begin{pmatrix} -\frac{1}{n-1} & -\frac{1}{n} & \cdots & -\frac{1}{n} \\ \frac{n-1}{n} & -\frac{1}{n} & \cdots & -\frac{1}{n} \\ \vdots & \vdots & \ddots & \vdots \\ -\frac{1}{n} & -\frac{1}{n} & \cdots & \frac{n-1}{n} \end{pmatrix}$$

and $v = 0$. Consequently, the region $\Omega_{s,n}^-$ where $\Xi_n(s; a_1, \dots, a_n) < 0$, plotted with respect to the variables $\log k_2, \dots, \log k_n$ is a convex and connected region containing the point $\log k_2 = \dots = \log k_n = 0$. Reducing the number of variables by setting some of the variables k_i to be equal to a constant or setting $k_i = k_j$ for some pairs of $i \neq j$ are tantamount to restricting the variables $(\log k_2, \dots, \log k_n)$ to the intersections of hyperplanes in $\mathbb{R}^{n-1}$. Therefore plotting the region $\Omega_{s,n}^-$ with respect to the remaining $\log k_i$ variables, the new region is still convex and connected. In [18], we have plotted the regions where $\Xi_3(s; a_1, a_2, a_3) < 0$ with respect to the variables $\log k_2$ and $\log k_3$, and the regions where $\Xi_4(s; a_1, a_2, a_3, a_4) < 0$ with respect to the variables $\log k_2, \log k_3$, with $k_4 = 1$ and $k_4 = 3$, for some values of s . The graphs show that these regions are indeed convex and connected. As a matter of fact, when the regions $\Omega_{s,n}^-$ are plotted using computer softwares, we can only determine the regions in finite domains of the $\log k$ variables. Our convexity and connectivity results guarantee that there does not exist region of $\Omega_{s,n}^-$ outside the finite domain we consider.

Finally, notice that the map $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n \mapsto (\log a_1, \dots, \log a_n) \in \mathbb{R}^n$ is continuous. On the other hand, we have shown that the intersection of the region $\Omega_{s,n}^-$ where $\Xi_n(s; a_1, \dots, a_n) < 0$ with any hypersurface in $(\mathbb{R}^+)^n$ of the form $\prod_{i=1}^n a_i = a$ a constant, contains the point $a_1 = \dots = a_n = a^{\frac{1}{n}}$. These allow one to conclude that the region $\Omega_{s,n}^-$, as a region of $(a_1, \dots, a_n) \in (\mathbb{R}^+)^n$, is a connected region that contains the ray $a_1 = \dots = a_n$.

5. CONCLUDING REMARKS

This paper is motivated by our recent work [18], in which we need to determine the conditions for which symmetry breaking occurs in the $\lambda\phi^4$ -interacting fractional Klein-Gordon field theory on a toroidal spacetime $T^n \times \mathbb{R}^N$. The results obtained in this work are used in [18]. Since the Epstein zeta function $Z_n(s; a_1, \dots, a_n)$ always appears when one studies field theories on toroidal manifolds or rectangular cavities, the results presented here may have potential applications for other works along these directions. For physics applications, one is usually interested in simple toroidal manifolds of the form $T^n \times \mathbb{R}^N$, which can be considered as quotients of $\mathbb{R}^n$ by rectangular lattices. This explains why we consider Epstein zeta function of the form $Z_n(s; a_1, \dots, a_n)$, $a \in (\mathbb{R}^+)^n$ instead of the general Epstein zeta function $Z_n(A; s)$ with A any positive definite symmetric matrix. An advantage of this simplification is that we can determine the minimum of $Z_n(s; a_1, \dots, a_n)$, $a \in (\mathbb{R}^+)^n$, $\prod_{i=1}^n a_i = 1$ for all $s \in \mathbb{R} \setminus \{0, n/2\}$ and all $n \in \mathbb{N}$. In contrast, for general Epstein zeta function $Z_n(A; s)$ with $\det A = 1$, only some local minima have been determined for $n = 2, 3, 4, 5, 6, 7, 8, 24$ with s in some range of $\mathbb{R}$ [15–38]. In fact, for all these known cases, the minimum of $Z_n(s; a_1, \dots, a_n)$, $a \in (\mathbb{R}^+)^n$, $\prod_{i=1}^n a_i = 1$ is no longer a local minimum in the larger domain where $A \in \{\text{positive definite symmetric } n \times n \text{ matrices}\}$. In the attempt to

search for the correct spacetime model, it might turn out that the general toroidal manifold $\mathbb{R}^n/L$, where L is any lattice in $\mathbb{R}^n$, will be of importance to physics. In that case, we need to extend the work of this paper to general Epstein zeta function $Z_n(A; s)$. It is hoped that the methods and results in this paper will be useful for the study of this general problem.

Our result about the convexity of the Epstein zeta function $Z_n(s; a_1, \dots, a_n)$, as a function of $\log a_1, \dots, \log a_n$ with $\log a_1 + \dots + \log a_n$ fixed, seems to be new. This result is important for determining the regions where $Z_n(s; a_1, \dots, a_n)$ is positive or negative, as discussed at the end of section 4. As a matter of fact, we have obtained a more general result regarding the convexity of functions $\mathcal{F}(x_1, \dots, x_n)$ of the form $\mathcal{F}(x_1, \dots, x_n) = \prod_{i=1}^n f(x_i)$, where $f(x)$ is a positive function that has continuous second derivative. This result may have applications in other areas of mathematics.

Acknowledgement The authors would like to thank Malaysian Academy of Sciences, Ministry of Science, Technology and Innovation for funding this project under the Scientific Advancement Fund Allocation (SAGA) Ref. No P96c.

REFERENCES

- [1] P. Epstein, *Zur Theorie allgemeiner Zetafunktionen*, Math. Ann. **56**, 615–644 (1903).
- [2] P. Epstein, *Zur Theorie allgemeiner Zetafunktionen II*, Math. Ann. **65**, 205–216 (1907).
- [3] C. L. Siegel, *Contributions to the theory of the Dirichlet L-series and the Epstein zeta-functions*, Ann. of Math. (2) **44**, 143–172 (1943).
- [4] M. Koecher, *Über Dirichlet-Reihen mit Funktionalgleichung*, J. Reine Angew. Math. **192**, 1–23 (1953).
- [5] C. L. Siegel, *A generalization of the Epstein zeta function*, J. Indian Math. Soc. (N.S.) **20**, 1–10 (1956).
- [6] A. Selberg, *A new type of zeta function connected with quadratic forms*, Report of the Inst. in Theory of Numbers, U. of Colorado, Boulder, Colo., 207–210 (1959).
- [7] A. Terras, *A generalization of Epstein's zeta function*, Nagoya Math. J. **42**, 173–188 (1971).
- [8] S. Chowla, *On an unsuspected real zero of Epstein's zeta function*, Proc. Nat. Inst. Sci. India **13**, no. 4, 1 (1947).
- [9] E. Elizalde, S. D. Odintsov, A. Romeo, A. A. Bytsenko, and S. Zerbini, *Zeta regularization techniques with applications*, (World Scientific Publishing Co. Inc., River Edge, NJ, 1994).
- [10] Emilio Elizalde, *Ten physical applications of spectral zeta functions*, (Springer-Verlag, Berlin, 1995).
- [11] K. Kirsten, *Spectral functions in mathematics and physics*, (Chapman & Hall/ CRC, Boca Raton, FL, 2002).
- [12] Jan Ambjørn and S. Wolfram, *Properties of the vacuum. I. Mechanical and thermodynamic*, Ann. Physics **147**, 1–32 (1983).
- [13] D. J. Toms, *Symmetry breaking and mass generation by space-time topology*, Phys. Rev. D **21**, 2805–2817 (1980).
- [14] G. Denardo and E. Spallucci, *Dynamical mass generation in $S^1 \times R^3$* , Nucl. Phys. B **169**, 514–526 (1980).
- [15] A. Actor, *Topological generation of gauge field mass by toroidal spacetime*, Class. Quantum Grav. **7**, 663–683 (1980).
- [16] K. Kirsten, *Topological gauge field mass generation by toroidal spacetime*, J. Phys. A: Math. Gen. **26**, 2421–2435 (1993).
- [17] E. Elizalde and K. Kirsten, *Topological symmetry breaking in self-interacting theories on toroidal space-time*, J. Math. Phys. **35**, 1260–1273 (1994).

- [18] S. C. Lim and L. P. Teo, *Topological symmetry breaking of self-interacting fractional Klein–Gordon field on toroidal spacetime*, submitted, can be downloaded from <http://pesona.mmu.edu.my/~lpteo>.
- [19] R. A. Rankin, *A minimum problem for the Epstein zeta-function*, Proc. Glasgow Math. Assoc. **1**, 149–158 (1953).
- [20] J. W. S. Cassels, *On a problem of Rankin about Epstein zeta function*, Proc. Glasg. Math. Assoc. **4**, 73–80 (1959); **6**, 116 (1963).
- [21] V. Ennola, *On a problem about the Epstein zeta-function*, Proc. Cambridge Philos. Soc. **60**, 855–875 (1964).
- [22] P. H. Diananda, *Notes on two lemmas concerning the Epstein zeta function*, Proc. Glasg. Math. Assoc. **6**, 202–204 (1964).
- [23] S. S. Ryskov, *On the question of the final ζ -optimality of lattices that yield the densest packing of n -dimensional balls*, Sibirsk. Mat. Ž. **14**, 1065–1075 (1973).
- [24] S. Š. Šušbaev, *An estimate of the origin of the ray of extremality of the principal perfect form in four variables*, Dokl. Akad. Nauk UzSSR, no. 9, 15–16 (1976).
- [25] S. Š. Šušbaev, *An estimate of the origin of the ray of extremality of the principal perfect form in five variables*, Voprosy Vyčisl. i Prikl. Mat. (Tashkent), no. 46, 3–8 (1977).
- [26] S. Š. Šušbaev, *The Rankin-Sobolev problem on extrema of Epstein's zeta function. The origin of the ray of extremality of the principal perfect form*, Izv. Akad. Nauk UzSSR Ser. Fiz.-Mat. Nauk, no. 1, 33–39 (1978).
- [27] K. M. Ėndibaev, *The extremal of Epstein's zeta function from four variables*, Voprosy Vychisl. i Prikl. Mat. (Tashkent), no. 51, 178–190 (1978).
- [28] A. Terras, *The minima of quadratic forms and the behavior of Epstein and Dedekind zeta functions*, J. Number Theory **12**, no. 2, 258–272 (1980).
- [29] K. M. Ėndibaev, *An extremal problem for the four-dimensional Epstein ζ -function*, Dokl. Akad. Nauk UzSSR, no. 4, 18–21 (1978).
- [30] K. L. Fields, *Locally minimal Epstein zeta functions*, Matematika **27**, no. 1, 17–24 (1980).
- [31] Z. D. Lomakina and S. S. Ryshkov, *One-dimensional and two-dimensional faces of the polyhedron $\mu(5)$* , Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **151**, no. 9, 95–103 (1986).
- [32] S. Sh. Shushbaev, *On the Rankin-Sobolev problem on extrema of Epstein zeta-functions and on a transcendental equation*, Voprosy Vychisl. i Prikl. Mat. (Tashkent), no. 82, 108–113 (1987).
- [33] S. Sh. Shushbaev, *Local minima of the Epstein zeta function*, Mat. Zametki **45**, no. 1, 123–131 (1989).
- [34] S. Sh. Shushbaev, *On the Rankin-Sobolev problem on extrema of the Epstein zeta-function. An estimate for the origin of a ray of extremality of the second perfect Voronoï form*, Sibirsk. Mat. Zh. **31**, no. 5, 157–163 (1990).
- [35] S. Sh. Shushbaev, *Two local minima of the six-dimensional Epstein zeta-function*, Izv. Vyssh. Uchebn. Zaved. Mat., no. 5, 78–83 (1990).
- [36] S. Sh. Shushbaev, *Rays of extremality: eventually extremal forms in seven variables*, Sibirsk. Mat. Zh. **35**, no. 6, 1397–1400 (1994).
- [37] E. V. Orlovskaya, *The minimum for a theta function of three variables and the solution of the Rankin-Sobolev problem in three-dimensional space*, Vestnik S.-Peterburg. Univ. Mat. Mekh. Astronom. **149**, no. 3, 23–27 (1994).
- [38] E. V. Orlovskaya, *The minimum for the theta-function of three variables and the solution of the Rankin-Sobolev problem in a three-dimensional space*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **211**, no. 3, 150–157 (1994).
- [39] W. Jenkner, *Sur les fonctions zêta attachées aux classes de rayon*, J. Théor. Nombres Bordeaux **7**, no. 1, 1–14 (1995).
- [40] S. Sh. Shushbaev and G. A. Kalybaeva, *On a perfect form that is not a final-extremal form*, Uzbek. Mat. Zh., no. 2, 80–86 (2002).
- [41] J. Steuding, *On the zero-distribution of Epstein zeta-functions*, Math. Ann. **333**, no. 3, 689–697 (2005).
- [42] P. Sarnak and A. Strömbergsson, *Minima of Epstein's zeta function and heights of flat tori*, Invent. Math. **165**, no. 1, 115–151 (2006).
- [43] S. Chowla and A. Selberg, *On Epstein's zeta function. I*, Proc. Nat. Acad. Sci. U. S. A. **35**, 371–374 (1949).

- [44] A. Selberg and S. Chowla, *On Epstein's zeta-function*, J. Reine Angew. Math. **227**, 86–110 (1967).
- [45] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*, (Academic Press Inc., San Diego, CA, 2000).
- [46] A. Fujii, *A note on the values of the Epstein zeta functions at the critical points*, Comment. Math. Univ. St. Pauli **49**, no. 2, 195–225 (2000).

E-mail address: ¹`sc1im@mmu.edu.my`

E-mail address: ²`lp1teo@mmu.edu.my`